

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 7/50, A47K 7/03		A1	(11) International Publication Number: WO 00/07561
			(43) International Publication Date: 17 February 2000 (17.02.00)
(21) International Application Number: PCT/EP99/05030 (22) International Filing Date: 14 July 1999 (14.07.99) (30) Priority Data: 09/130,981 7 August 1998 (07.08.98) US (71) Applicant (for AE AU BB CA CY GB GD GII GM IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZA ZW only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AE AU BB CA CY GB GD GH GM IE IL IN KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZA ZW): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). (71) Applicant (for IN only): HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Mumbai 400 020, Maharashtra (IN).		(72) Inventors: FARRELL, Linda; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). SLAVTCHEFF, Craig, Stephen; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). ZNAIDEN, Alexander, Paul; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). VINSKI, Paul; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). (74) Agent: ROTS, Maria, Johanna, Francisca; Unilever PLC, Patent Department, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: COSMETIC EFFERVESCENT CLEANSING PILLOW			
(57) Abstract			
<p>A wiping article is provided which includes an effervescent cleanser composition held within a pouch formed from a first and second substrate sheet. At least one of the substrate sheets must be water permeable. The effervescent composition is an intimate mixture of an acid material such as citric acid and an alkaline material such as sodium bicarbonate. Water contact causes the combination to effervesce. A dry surfactant such as sodium cocoyl isethionate in contact with the water and effervescing carbon dioxide results in a highly pleasant sudsing system. Skin benefit agents may be included within the composition. The effervescent action may improve deposition of the skin benefit agents onto the skin.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

COSMETIC EFFERVESCENT CLEANSING PILLOW

The invention concerns an effervescent foaming wipe article for body cleansing that imparts a pleasant sensory feel to a user's skin.

Classically the process of cleansing skin or other articles has employed a surfactant composition. Sometimes an implement has joined the composition. Implements such as sachets serve a multi-purpose. One function is as a delivery package for the surfactant. Sachets may also assist in generating foam. They also function as an abrasive assisting in the cleansing function.

An early example of cleansing pad technology is found in U.S. Patent 1,808,834 (Busch Sr.). A fabric pouch is disclosed surrounding a cleansing composition mainly consisting of calcium and sodium carbonate.

U.S. Patent 4,234,442 (Cornelissens) describes a sachet which can consist of a water permeable material filled with an acidic and an alkaline constituent. Adipic, succinic and glutaric acids exemplify the acidic constituent. Sodium bicarbonate and carbonate represent the alkaline ingredient.

25

U.S. Patent 4,272,393 (Gergely) describes a cleaning article formed of a porous flexible substrate, especially a cellulosic paper, impregnated with detergent and a gas-generating system. The latter is formed by separating an acidic component such as citric acid from a basic component

30

- 2 -

such as sodium carbonate in two separate areas of the substrate.

U.S. Patent 4,515,703 (Haq), U.S. Patent 4,600,620 (Lloyd et al.) and U.S. Patent 4,603,069 (Haq et al.) all describe
5 wiping articles impregnated with surfactant. These do not contain any effervescent ingredients.

WO 97/43366 (Askew et al.) reports an effervescent system to
10 improve dispensability of granular laundry detergent powders into wash water of automatic washing machines. Citric acid and bicarbonate combinations are employed to generate effervescence.

15 It is an object of the present invention to provide a cleansing cloth containing an effervescent system activated by contact with water.

Another object of the present invention is to provide a
20 cleansing cloth containing an effervescent system activated by contact with water to generate carbon dioxide which expands the lathering of cleanser components of the cloth.

Still a further object of the invention is to provide a
25 cleansing cloth which imparts a pleasant sensory feel to skin during and after use.

According to a first aspect, there is provided a cosmetic
30 article is provided for cleansing body surfaces, the article including:

- 3 -

a pouch formed of first and second sheets, at least one being water permeable, the first and second sheets defining a pouch between them, the pouch being sealed along all its perimeter; and

5 an effervescent cleanser composition in the form of an anhydrous dry solid being positioned within the pouch, the composition including:

- (i) from about 1 to about 80% of an alkaline material;
- (ii) from about 0.5 to about 80% of an acid material; and
- 10 (iii) from about 0.1 to about 30% of a solid surfactant.

Also provided is a method for cleansing skin involving wetting with water a cosmetic cleansing article, generating foam from the article and wiping skin surfaces with the
15 wetted article, the article being the pouch with effervescent cleanser composition described above.

Cosmetic wiping articles of the present invention when contacted with water billow to many times (e.g. typically
20 more than 10, but often more than 40 times) their dry size when activated by water. The effervescent cleansing system exudes copious amounts of lather. A plumped "pillow" arises from the effervescent action. By careful control of the acidic and alkaline components, a squeaky clean rinsed
25 feeling is felt on a user's skin.

A first essential component of compositions within the pouch is that of an acidic material. Suitable for this purpose are any acids present in dry solid form. Especially
30 appropriate are C₂-C₂₀ organic mono- and poly- carboxylic acids and especially alpha- and beta- hydroxycarboxylic

- 4 -

acids; C₂-C₂₀ organophosphorus acids such as phytic acid; C₂-C₂₀ organosulfur acids such as toluene sulfonic acid; and peroxides such as hydrogen peroxide. Typical hydroxycarboxylic acids include adipic, glutaric, succinic, tartaric, malic, maleic, lactic, salicylic and citric acids as well as acid forming lactones such as gluconolactone and glucarolactone. Most preferred is citric acid.

Also suitable as acid material may be encapsulated acids. Typical encapsulating material may include water soluble synthetic or natural polymers such as polyacrylates (e.g. encapsulating polyacrylic acid), cellulosic gums, polyurethane and polyoxyalkylene polymers. By the term "acid" is meant any substance which when dissolved in deionized water at 1% concentration will have a pH of less than 7, preferably less than 6.5, optimally less than 5. These acids preferably at 25°C are in solid form, i.e. having melting points no less than 25°C. Concentrations of the acid should range from about 0.5 to about 80%, preferably from about 10 to about 65%, optimally from about 20 to about 45% by weight of the total composition.

A second essential component of compositions within the pouch is that of an alkaline material. The alkaline material is a substance which can generate a gas such as carbon dioxide, nitrogen or oxygen, i.e. effervesce, when contacted with water and the acidic material. Suitable alkaline materials are anhydrous salts of carbonates and bicarbonates, alkaline peroxides (e.g. sodium perborate and sodium percarbonate) and azides (e.g. sodium azide). Preferably the alkaline material is sodium or potassium

- 5 -

bicarbonate. Amounts of the alkaline material may range from about 1 to about 80%, preferably from about 5 to about 49%, more preferably from about 15 to about 40%, optimally from about 25 to about 35% by weight of the total composition.

By the term "anhydrous" is meant the presence of no more than 5%, preferably no more than 3.5% and optimally no more than 1% water by weight of the total composition. Water of hydration is not considered to be water for purposes of the anhydrous definition. However, it is preferred to minimize, and preferably to eliminate any water of hydration.

Advantageously the combined amount of acidic and alkaline materials will be at least about 1.5%, preferably from about 40 to about 95%, optimally from about 60 to about 80% by weight of the total composition.

A third necessary component of compositions according to the present invention is that of a dry surfactant, preferably a dry surfactant which is solid at 20°C. Most suitable for the present invention is sodium cocoyl isethionate. Other useful surfactants include sodium methyl cocoyl taurate and sodium lauryl sulfate. Surfactants may be of the anionic, cationic, nonionic, amphoteric, zwitterionic varieties and combinations thereof. Amounts of the dry surfactant may range from about 0.1 to about 30%, preferably from about 1 to about 30%, optimally from about 8 to about 15% by weight of the total composition.

- 6 -

A variety of skin benefit agents may be included to improve afterfeel properties. Advantageously these substances will be available as anhydrous dry powders. Alternatively these substances may be liquids deposited upon or into a powdered
5 substrate (e.g. sodium bicarbonate or zeolite) to achieve a resultant dry flowing powder.

Within the skin benefit agent scope are several suitable categories of materials. These include emollients, anti-
10 aging actives, antibacterials and fungicides, skin lighteners, sunscreens and combinations thereof. Amounts of the skin benefit agents may range from about 0.001 to about 30%, preferably from about 0.1 to about 20%, more preferably from about 0.5 to about 10%, optimally between about 1 and
15 about 5% by weight of the total composition.

Emollients may be in the form of natural or synthetic esters, silicone oils, hydrocarbons, starches, fatty acids and mixtures thereof. Typically the emollient may range in
20 concentration from about 0.1 to about 35% by weight of the total composition.

Suitable silicone oils may be divided into the volatile and nonvolatile variety. The term "volatile" as used herein
25 refers to those materials which have a measurable vapor pressure at ambient temperature. Volatile silicone oils are preferably chosen from cyclic or linear polydimethylsiloxanes containing from 3 to 9, preferably from 4 to 5, silicon atoms.

- 7 -

Linear volatile silicone materials generally have viscosities less than about 5 centistokes at 25°C while cyclic materials typically have viscosities of less than about 10 centistokes.

5

Nonvolatile silicone oils useful as an emollient material include polyalkyl siloxanes, polyalkylaryl siloxanes and polyether siloxane copolymers. The essentially non-volatile polyalkyl siloxanes useful herein include, for example, polydimethyl siloxanes with viscosities of from about 5 to about 100,000 centistokes at 25°C. Among the preferred non-volatile emollients useful in the present compositions are the polydimethyl siloxanes having viscosities from about 10 to about 400 centistokes at 25°C.

15

Among the suitable ester emollients are:

(1) Alkenyl or alkyl esters of fatty acids having 10 to 20 carbon atoms. Examples thereof include isoarachidyl neopentanoate, isononyl isonanonoate, oleyl myristate, oleyl stearate, and oleyl oleate.

20

(2) Ether-esters such as fatty acid esters of ethoxylated fatty alcohols.

25

(3) Polyhydric alcohol esters. Ethylene glycol mono and di-fatty acid ester, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol (200-6000) mono- and di-fatty acid esters, polypropylene glycol 2000 monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty

30

- 8 -

acid esters, polyglycerol poly-fatty esters, ethoxylated glyceryl monostearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters are satisfactory polyhydric alcohol esters.

(4) Wax esters such as beeswax, spermaceti, myristyl myristate, stearyl stearate and arachidyl behenate.

10

(5) Sterols esters, of which cholesterol fatty acid esters are examples thereof.

(6) Triglycerides such as sunflower seed oil, maleated sunflower seed oil, borage seed oil and safflower oil.

15

Hydrocarbons suitable as emollients include petrolatum, mineral oil, isoparaffins and hydrocarbon waxes such as polyethylene.

20

Starches are also suitable emollients. Typical of this class is tapioca and arabinogalactan.

Fatty acids may also be suitable as emollients. The fatty acids normally have from 10 to 30 carbon atoms.

25

Illustrative of this category are pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic, oleic, linoleic, ricinoleic, arachidic, behenic and erucic acids.

30

- 9 -

Anti-aging actives are also useful as skin benefit agents. Included within this category are vitamins, retinoids and combinations thereof. Amounts of these materials may range from about 0.001 to about 20% by weight of the total

5 composition. Suitable vitamins include ascorbic acid, Vitamin B₆, Vitamin B₁₂, tocopherol as well as salts and C₁-C₂₀ esters thereof. Suitable retinoids include retinoic acid as well as its C₁-C₂₂ esters and salts, retinol and C₁-C₂₂ fatty esters of retinol including retinyl linoleate.

10

Another class of anti-aging actives are the alpha- and beta-hydroxycarboxylic acids and salts thereof. Representative of this group are glycolic acid, lactic acid, malic acid, hydroxyoctanoic acid and mixtures of these as well as their

15 salts. Suitable salts are the alkalimetal, ammonium and C₁-C₁₀ alkanol ammonium salts.

Antibacterials and fungicidals may also be included as skin benefit agents. Representative of these categories are

20 triclosan, tricloban, hexetidene, chlorhexadene, gluconates, zinc salts (e.g. zinc citrate and zinc phenolsulfonate) and combinations thereof.

Skin lighteners may also be included under the skin benefit

25 agents. Typical of this category are niacinamide, kojic acid, arbutin, vanillin, ferulic acid and esters thereof, resorcinol, hydroquinone, placental extract and combinations thereof.

30 Sunscreens may also be included as skin benefit agents. Particularly preferred are such materials as ethylhexyl p-

- 10 -

methoxycinnamate, available as Parsol® MCX, and benzophenone-3, also known as Oxybenzone. Inorganic sunscreen actives may be employed such as microfine titanium dioxide, polyethylene and various other polymers. Amounts
5 of the sunscreen agents will generally range from 0.1 to 30%, preferably from 2 to 20%, optimally from 4 to 10% by weight.

Adjunct functional agents may also be incorporated into
10 compositions of the present invention. These include electrolytes, thickeners and mixtures thereof. Amounts of these substances may range from about 0.1 to about 20%, preferably from about 0.3 to about 10%, optimally between about 0.5 and about 5% by weight of the total composition.

15 Electrolytes may be selected from alkali, alkaline earth or ammonium salts of phosphates, silicates, halides, sulphates and mixtures thereof. Typical phosphates are potassium polymetaphosphate, sodium tripolyphosphate, sodium
20 tetrapyrophosphate, sodium or potassium pyrophosphate and sodium hexametaphosphate. Most preferred is potassium polymetaphosphate available as Lipothix 100B® which is a 70:30 mixture of potassium polymetaphosphate and sodium bicarbonate, available from Lipo Chemicals, Inc., Paterson,
25 New Jersey. Preferred sulphates are the magnesium sulphates.

Thickeners which may improve afterfeel properties on skin include inorganic or organic substances. A particularly
30 preferred inorganic thickener is sodium magnesium silicate commercially available as Optigel SH®. Organic thickeners

- 11 -

include alginic acid as well as sodium and calcium alginates, sodium carboxymethyl cellulose, hydroxypropyl methylcellulose, hydroxypropyl cellulose, hydroxyethyl cellulose and combinations thereof.

5

Most preferred is alginic acid, which is commercially available as Kelacid® from Sud-Chemie Rheologicals, Louisville, Kentucky. Alginic acid is highly effective at removing the slimy feel associated with deposits of alkaline material which are not fully rinsed away from the skin. Amounts of the thickener may range from about 0.1 to about 20%.

Polysaccharides useful in this invention are dry solid anhydrous substances such as sorbitol, sugars, (such as trehalose) starches, modified starches (e.g. aluminum octenyl succinate) and mixtures thereof. Most preferred is sorbitol.

Deposition aids may also be incorporated in compositions of the present invention. These assist in depositing skin benefit agents onto the skin surface. Particularly effective are cationic monomers and polymers for this purpose. Illustrative are the following:

25

Lauryltrimethylammonium chloride (Laurtrimonium chloride);

Stearyltri(2-hydroxyethyl) ammonium chloride (Quaternium- 16);

30

Lauryldimethylbenzylammonium chloride (Lauralkonium chloride);

- 12 -

- Oleyldimethylbenzylammonium chloride (Olealkonium chloride);
- 5 Dilauryldimethylammonium chloride (Dilauryldimonium chloride);
- Cetyltrimethylbenzylammonium chloride (Cetalkonium chloride);
- 10 Dicetyltrimethylammonium chloride (Dicetyltrimonium chloride);
- Laurylpyridinium chloride (Laurylpyridinium chloride);
- 15 Cetylpyridinium chloride (Cetylpyridinium chloride);
- N-(soya alkyl)-N,N,N-trimethylammonium chloride (Soyatrimonium chloride);
- 20 Polydiallyldimethylammonium chloride (Polyquaternium-6);
- Diallyldimethylammonium salt copolymerized with acrylamide (Polyquaternium-7);
- 25 Guar hydroxypropyltrimonium Chloride (Guar hydroxypropyltrimonium chloride);
- 30 Copolymer of N-vinyl-pyrrolidone and N,N-dimethylaminoethylmethacrylate, quaternized with dimethylsulfate (Polyquaternium-11);
- 35 Copolymer of acrylamide and N,N-dimethylaminoethyl methacrylate, quaternized with dimethyl sulfate (Polyquaternium-5);
- Cationic hydroxyethylcellulosics (Polyquaternium-10);
- Cationic hydroxyethylcellulosics (Polyquaternium-24);
- 40 Cetyltrimethylammonium chloride (Cetrimonium chloride);
- Decyldimethyloctylammonium chloride (Quaternium-24);
- 45 Myristyltrimethylammonium chloride (Mytrimonium chloride);

- 13 -

- Polyoxyethylene (2)-cocomonium Chloride (PEG-2 Cocomonium chloride);
- 5 Methylbis(2-hydroxyethyl) cocoammonium chloride (PEG-2 Cocoyl Quaternium-4);
- Methylpolyoxyethylene-(15) cocoammonium chloride (PEG-15 Cocoyl Quaternium-4);
- 10 Methylbis(2-hydroxyethyl) octadecyl ammonium chloride (PEG-2 Stearyl Quaternium-4);
- 15 Methylpolyoxyethylene-(15) octadecylammonium chloride (PEG-15 Stearyl Quaternium-4);
- Methylbis(2-hydroxyethyl)-oleylammonium chloride (PEG-2 Oleyl Quaternium-4);
- 20 Methylpolyoxyethylene-(15) oleylammonium chloride (PEG-15 Oleyl Quaternium-4);
- 25 The names in parenthesis are given by the Cosmetic, Toiletry and Fragrance Association, Inc. in the CTFA Cosmetic Ingredient Dictionary. Most preferred for purposes of this invention are cationic guar gums such as Jaguar C13S®, which is guar hydroxypropyltrimonium chloride. Amounts of the
- 30 deposition aid may range from about 0.01 to about 1%, preferably from about 0.05 to about 0.5%, optimally from about 0.1 to about 0.3% by weight.

Advantageously an emotive agent such as a fragrance and/or

35 botanical extract are included with the effervescent cleansing composition. Fragrances and botanicals are often liquids. For this reason it is necessary to uniformly distribute and allow absorption of liquid components into the solid powder. One method of best achieving this is to

- 14 -

spray these liquids onto the solids. Amounts of the fragrance and/or botanicals combined may be at levels from about 0.1 to about 3%, preferably from 0.5 to 2%, optimally from 0.8 to 1.5% by weight of the total composition.

5

The term "fragrance" is defined as a mixture of odoriferous components, optionally mixed with a suitable solvent diluent or carrier, which is employed to impart a desired odor.

Particular preferred odoriferous components are cyclic and
10 acyclic terpenes and terpenoids. These materials are based upon isoprene repeating units. Examples include alpha and beta pinene, myrcene, geranyl alcohol and acetate, camphene, dl-limonene, alpha and beta phellandrene, tricyclene, terpinolene, allocimmane, geraniol, nerol, linanool,
15 dihydrolinanool, citral, ionone, methyl ionone, citronellol, citronellal, alpha terpineol, beta terpineol, alpha fenchol, borneol, isoborneol, camphor, terpinen-1-ol, terpin-4-ol, dihydroterpineol, methyl chavicol, anethole, 1,4 and 1,8 cineole, geranyl nitrile, isobornyl acetate, linalyl
20 acetate, caryophyllene, alpha cedrene, guaiol, patchouli alcohol, alpha and beta santalol and mixtures thereof.

Botanicals of particular use in the present invention include yarrow, chamomile, jasmine, lavender, horse
25 chestnut, sage, thyme, yucca, coltsfoot and mixtures thereof.

Preservatives can desirably be incorporated into the cosmetic compositions of this invention to protect against
30 the growth of potentially harmful microorganisms. Suitable traditional preservatives for compositions of this invention

- 15 -

are alkyl esters of para-hydroxybenzoic acids. Other preservatives which have more recently come into use include hydantoin derivatives, propionate salts, and a variety of quaternary ammonium compounds. Cosmetic chemists are familiar with appropriate preservatives and routinely choose them to satisfy the preservative challenge test and to provide product stability.

Particularly preferred preservatives are phenoxyethanol, methyl paraben, propyl paraben, imidazolidinyl urea, sodium dehydroacetate and benzyl alcohol. The preservatives should be selected having regard for the use of the composition and possible incompatibilities between the preservatives and other ingredients in the emulsion. Preservatives are preferably employed in amounts ranging from 0.01% to 2% by weight of the composition.

Natural vegetable materials from renewable resources are often desirable in cosmetic compositions. For instance, cosmetic compositions of the present invention may include beta-glucan derived from oats, commercially available under the trademark Microat SF from Nurture Inc., Missoula, Mont.

Colorants may also be included in compositions of the present invention. These substances may range from about 0.05 to about 5%, preferably between 0.1 and 3% by weight.

Effervescent cleansing compositions of this invention will be placed within a pouch formed between a first and second flexible substrate sheet, preferably at least one of these being a flexible sheet. At least one of the sheets must be

- 16 -

water permeable, most preferably both sheets should have water permeability. For definitional purposes, the first and second sheets can be folded-over panels of a single unitary sheet.

5

Suitable materials for forming sheets may be rayon, polyester, polyethylene, polypropylene, cotton or any combination thereof. These sheets may be woven or non-woven. Most preferred is a non-woven rayon. Cellulosic paper fiber substrates are best not employed because of their insufficient wet-strength, although they may be blended with other fibers referenced above; it is important that the substrate sheets are not readily torn open through consumer rubbing of the article. Unlike laundry sachet articles, pouches of the present invention should not rupture to allow dispersion of their granular contents into wash water. Rather it is intended for all cleanser composition components to exit by dissolution through the permeable walls of the pouch.

20

Skin surfaces against which articles of the present invention are useful include face, body, scalp, axilla and even legs/feet. When the article is a foot cleanser, it would be advantageous for the pouch on one of its sides to be coarse while the second of the sheets may be soft and gentle. An abrasive non-woven flexible sheet in a foot cleanser product is useful for rubbing against calluses while the second sheet of the pouch remains smooth.

25

Articles according to the present invention may be formed in the following manner. Constituents of the effervescent

30

- 17 -

cleansing composition are placed into a dry mill or similar apparatus, and blended until a uniformly distributed powder results. Thereafter, fragrance/botanical component as a second phase is sprayed into the dry mill with concurrent
5 agitation of the powdered composition.

A continuous roll of first substrate sheet is unwound from a source roll over a moving conveyer belt. The effervescent cleansing composition is placed into a hopper positioned
10 over the conveyer belt. A discrete charge of powdered composition is regularly placed on the first substrate sheet at a location directly under a nozzle of the hopper. A second substrate sheet is then in register placed over that of the loaded first substrate sheet. At this point all four
15 corners defining a rectangle or square are sealed in register trapping the effervescent cleansing composition within. Cutters then separate one sealed section from another thereby forming the wiping article. One or more of the wiping articles are then packaged within a moisture
20 impermeable outer package such as a laminated foil bag to prevent activation of the effervescent system during storage.

Ultrasonic welding may be employed as an alternative to
25 heat-sealing of the first and second substrates together. Thread stitching, glue application or other closure mechanisms may also be utilized.

Except in the operating and comparative examples, or where
30 otherwise explicitly indicated, all numbers in this

- 18 -

description indicating amounts of material are to be understood as modified by the word "about".

The term "comprising" is meant not to be limiting to any subsequently stated elements but rather to encompass non-specified elements of major or minor functional importance. In other words the listed steps, elements or options need not be exhaustive.

The following examples will more fully illustrate embodiments of this invention. All parts, percentages and proportions referred to herein and in the appended claims are by weight unless otherwise indicated.

EXAMPLE 1

An effervescent cleansing composition was prepared according to the formulation reported in Table I. Phase A was dry blended in a high speed shearing mixer. Fragrance was then sprayed onto the resultant powder as a Phase B. Three grams of the resultant powder were then placed into a 5.1cm by 7.6 cm (two inch by three inch) pouch formed of non-woven rayon. All sides were closed by double stitching with thread.

- 19 -

TABLE I

INGREDIENT	WEIGHT %
PHASE A	
Sodium Bicarbonate	34.5
Citric Acid (Anhydrous)	40.4
Sodium Cocoyl Isethionate (Powder)	11.6
Sodium Sesquicarbonate	5.0
Lipothix 100B® (Potassium Polymetaphosphate/Bicarbonate 70:30)	0.5
Optigel SH® (Sodium Magnesium Silicate)	1.0
Kelacid® (Alginic Acid)	1.0
Sorbitol	5.0
PHASE B	
Fragrance	1.0

EXAMPLE 2

5

Another effervescent cleansing composition was prepared according to the formulation reported in Table II.

TABLE II

10

INGREDIENT	WEIGHT %
PHASE A	
Sodium Bicarbonate	32.3
Citric Acid (Anhydrous)	41.1
Sodium Cocyl Isethionate (Powder)	11.6
Sodium Sesquicarbonate	5.0
Lipothix 100B® (Potassium Polymetaphosphate/Bicarbonate 70:30)	0.5
Optigel SH® (Sodium Magnesium Silicate)	1.0
Kelacid® (Alginic Acid)	1.0
Sorbitol	5.0
Laracare A200® (Arabinogalactan)	1.0
Ascorbic Acid	0.5
PHASE B	
Fragrance	1.0

EXAMPLE 3

A face cleansing effervescent composition was prepared
5 according to the formulation reported in Table III.

TABLE III

INGREDIENT	WEIGHT %
PHASE A	
Sodium Bicarbonate	33.6
Citric Acid (Anhydrous)	39.0
Sodium Cocyl Isethionate (Powder)	3.0
Sodium Methyl Cocoyl Taurate	6.0
Sodium Lauryl Sulfate	2.5
Sodium Sesquicarbonate	5.0
Lipothix 100B® (Potassium Polymetaphosphate/Bicarbonate 70:30)	0.5
Optigel SH® (Sodium Magnesium Silicate)	2.0
Tapioca	5.5
Methyl Gluceth 20-Benzate	2.0
Guar Hydroxypropyl Trimonium Chloride	0.25
PHASE B	
Fragrance	0.65

10 **EXAMPLE 4**

A still further effervescent cleansing composition according
to the present invention may be prepared according to the
formulation reported under Table IV. Phase A is prepared by
15 dry mixing of the ingredients in a high speed shear mixer.
Three grams of resultant powder are placed into a 5.1cm by
7.6 cm (two inch by three inch) pouch formed of non-woven
cotton polyester (50:50). The mesh size of the pouch walls
is sufficient to allow transfer of dissolved ingredients.

- 21 -

All sides of the pouch are welded by ultrasonic heat to insure against powder escaping from the pouch. TABLE IV

INGREDIENT	WEIGHT %
PHASE A	
Potassium Bicarbonate	29.5
Lactic Acid (Anhydrous)	45.4
Sodium Sulfosuccinate	11.6
Sodium Sesquicarbonate	5.0
Lipothix 100B® (Potassium Polymetaphosphate/Bicarbonate 70:30)	0.5
Optigel SH® (Sodium Magnesium Silicate)	1.0
Kelacid® (Alginic Acid)	1.0
Sorbitol	5.0
PHASE B	
Fragrance	1.0
Licorice Extract	0.1

5 EXAMPLE 5

Still another effervescent cleansing composition is prepared according to the formulation reported in Table V. The ingredients are dry blended in a high speed shearing mixer.

- 10 Fragrance and herbal extract are sprayed onto the powder and further blended to achieve homogeneity. Three grams of the resultant powder are placed into a 5.1cm by 7.6 cm (two inch by three inch) pouch formed of non-woven polypropylene. All sides are closed by convection heat sealing along the
- 15 perimeter thereof.

- 22 -

TABLE V

INGREDIENT	WEIGHT %
PHASE A	
Sodium Bicarbonate	29.5
Citraconic Acid (Anhydrous)	45.4
Methyl Glucamide	11.6
Sodium Sesquicarbonate	5.0
Lipothix 100B® (Potassium Polymetaphosphate/Bicarbonate 70:30)	0.5
Optigel SH® (Sodium Magnesium Silicate)	1.0
Kelacid® (Alginic Acid)	1.0
Sorbitol	5.0
PHASE B	
Fragrance	0.9
Yarrow	0.1

- 5 The foregoing description and examples illustrate selected embodiments of the present invention. In light thereof variations and modifications will be suggested to one skilled in the art, all of which are within the spirit and purview of this invention.

CLAIMS

1. A cosmetic article for cleansing body surfaces, the article comprising:

5 a pouch formed of first and second sheets, at least one being water permeable, the first and second sheets defining a pouch between them, the pouch being sealed along all its perimeter; and

an effervescent cleanser composition in the form
10 of an anhydrous dry solid being positioned within the pouch, the composition comprising:

- (i) from 1 to 80% of an alkaline material;
- (ii) from 0.5 to 80% of an acid material; and
- (iii) from 0.1 to 30% of a solid surfactant.

15

2. The article according to claim 1 further comprising from 0.01 to 30% by weight of a skin benefit agent selected from emollients, anti-aging actives, antibacterials and fungicides, skin lighteners, sunscreens and mixtures
20 thereof.

25

3. The article according to claim 2 wherein the anti-aging actives are selected from the group consisting essentially of vitamins, retinoids and mixtures thereof.

4. The article according to any of the preceding claims further comprising from 0.1 to 3% by weight of an emotive agent selected from fragrance, botanical and mixtures
thereof.

30

- 24 -

5. The article according to claim 4 wherein the emotive agent is incorporated into the composition by spray application onto one or more other components of the composition.

5

6. The article according to any of the preceding claims wherein the water permeable substrate sheet is a non-woven rayon.

10 7. The article according to any of the preceding claims wherein the surfactant is sodium cocoyl isethionate.

8. The article according to any of the preceding claims wherein the acid is citric acid.

15

9. The article according to any of the preceding claims wherein the composition further comprises from 0.01 to 1% by weight of a deposition aid which is a cationic monomer or polymer.

20

10. A method for cleansing skin comprising wetting with water a cosmetic cleansing article, generating foam from the article and wiping skin surfaces with the wetted article, the article comprising:

25

a pouch formed of first and second sheets, at least one being water permeable, the first and second sheets defining a pouch between them, the pouch being sealed along all its perimeter; and

30 an effervescent cleanser composition in the form of an anhydrous dry solid being positioned within the pouch, the composition comprising:

- 25 -

- (i) from 1 to 80% of an alkaline material;
- (ii) from 0.5 to 80% of an acid material; and
- (iii) from 0.1 to 30% of a solid surfactant.

INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/EP 99/05030

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K7/50 A47K7/03

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A47K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE 197 45 964 A (WELLA) 25 June 1998 (1998-06-25) page 2, line 65 -page 3, line 27; claims 1-9; example 4 ---	1-10
Y	GB 2 118 961 A (FIRMENICH) 9 November 1983 (1983-11-09) page 2, right-hand column, line 52 - line 61; claims 1-3 ---	1-10
Y	PATENT ABSTRACTS OF JAPAN vol. 11, no. 232 (C-437) '2679!, 29 June 1987 (1987-06-29) & JP 62 045519 A (KING KAGAKU), 27 February 1987 (1987-02-27) abstract --- -/--	1-10

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

1 November 1999

Date of mailing of the international search report

10/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Willekens, G

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 806 201 A (BRISTOL-MYERS SQUIBB) 12 November 1997 (1997-11-12) page 5, line 43 - line 47; claims 1-22; example 1 ---	1-10
A	EP 0 423 015 A (L'OREAL) 17 April 1991 (1991-04-17) page 4, line 23 - line 28; claims 1-21; example 3 ---	1-10
P,X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 1998-550915 XP002121154 & JP 10 245075 A (KAWAI), 14 September 1998 (1998-09-14) abstract ---	1-10
P,X	WO 98 42303 A (UNILEVER) 1 October 1998 (1998-10-01) claims 1-10 -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05030

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 19745964	A	25-06-1998	NONE	
GB 2118961	A	09-11-1983	FR 2525107 A	21-10-1983
			DE 3313744 A	20-10-1983
			DE 8311265 U	24-09-1987
			IT 1163199 B	08-04-1987
			JP 58188813 A	04-11-1983
JP 62045519	A	27-02-1987	JP 1456017 C	09-09-1988
			JP 63002935 B	21-01-1988
EP 806201	A	12-11-1997	US 5720949 A	24-02-1998
			CA 2202735 A	06-11-1997
EP 423015	A	17-04-1991	FR 2653013 A	19-04-1991
			AT 88886 T	15-05-1993
			AU 628673 B	17-09-1992
			AU 6454690 A	18-04-1991
			CA 2027483 A,C	14-04-1991
			DE 69001535 T	05-01-1994
			JP 3227910 A	08-10-1991
			JP 6025052 B	06-04-1994
			US 5100674 A	31-03-1992
JP 10245075	A	14-09-1998	NONE	
WO 9842303	A	01-10-1998	AU 6729298 A	20-10-1998
			AU 6830898 A	20-10-1998
			AU 7041798 A	20-10-1998
			WO 9842302 A	01-10-1998
			WO 9842304 A	01-10-1998
			US 5935596 A	10-08-1999